

Publikationsverzeichnis

Prof. Dr. Klaus G. Reymann

Originalarbeiten:

Reymann, K., Grincenko, Ju.V. und Svyrkov, V.B.: Aktivitätsänderungen hippocampaler Neuren beim instrumentellen Nahrungsverhalten. **Med. u. Sport, 1977, 17, 203-204**

Reymann, K., Svyrkov, V.B. und Grincenko, Ju.V.: Über die Beteiligung hippocampaler Neuren an den Teilschritten eines Nahrungsverhaltens. **Acta biol. med. germ., 1977, 36, 1107-1112**

Reymann, K., Walek, H. und Rost, R.: EEG-Leistungsspektrum des visuellen Kortex des Kaninchens bei Langzeitexperimenten. **Acta biol. med. germ., 1978, 37, 1454-1460**

Reymann, K. und Schmidt, D.: Beeinflussung der Herzfrequenz des Kaninchens durch Nahrungsdeprivation. **Acta biol. med. germ., 1979, 38, 691-694**

Reymann, K., Schmidt, D. und Haubenreiser, J.: Änderungen photisch evozierter Potentiale im Nahrungsverhalten von Kaninchen. 1. Abhängigkeit vom motivationellen Zustand. **Acta biol. med. germ., 1980, 39, 381-391**

Reymann, K. und Haschke, W.: Änderungen photisch evozierter Potentiale im Nahrungsverhalten von Kaninchen. 2. Abhängigkeit von der biologischen Relevanz des Signals. **Acta biol. med. germ., 1980, 39, 393-401**

Ott, T., Rüthrich, H., Reymann, K., Lindenau, L., and Matthies, H.: Direct evidence for the participation of changes in synaptic efficacy in the development of behavioral plasticity. In: *Neuronal Plasticity and Memory Formation*: G.A. Marsan and Matthies (eds.) **Raven Press, New York, 1982, 441-452**

Reymann, K., Rüthrich, H., Lindenau, L., Ott, T., and Matthies, H.: Monosynaptic activation of the hippocampus as a conditioned stimulus: Behavioral effects. **Physiol. Behav., 1982, 29, 1007-1012**

Reymann, K.G., and Ott, T.: Footshock-induced modification of a monosynaptic evoked potential in hippocampus. **Physiol. Behav., 1983, 31, 615-617**

Reymann, K.G., Pohle, W., Welde-Müller, P., and Ott, T.: Dopaminergic innervation of the hippocampus: Evidence for midbrain raphe neurons as the site of origin. **Biomed. Biochem. Acta, 1983, 42, 1239-1247**

Reymann, K.G., Chepkova, A.N., and Matthies, H.: Effects of deprolorphan, a casomorphin analog, on hippocampal CA1 field potentials in vitro. **Peptides, 1983, 4, 283-286**

Reymann, K.G., Chepkova, A.N., Schulzeck, K., and Ott, T.: Effects of β -casomorphin on dentate hippocampal field potentials in freely moving rats. **Biomed. Biochem. Acta**, 1985, 44, 749-754

Reymann, K.G., Malisch, R., Schulzeck, K., Brödemann, R., Ott, and Matthies H.: The duration of long-term potentiation in the CA1 region of the hippocampal slice preparation. **Brain Res. Bull.**, 1985, 15, 249-255

Reymann, K.G., Wulcko, S., Ott, T., and Matthies, H.: Opioid-receptor blockade reduces nose-poke self-stimulation derived from medial entorhinal cortex. **Pharmacol. Biochem. Behav.**, 1986, 24, 439-443

Thiemann, W., Malisch, R., and Reymann, K.G.: A new microcirculation chamber for inexpensive long-term investigations of nervous tissue in vitro. **Brain Res. Bull.**, 1986, 17, 1-4

Reymann, K.G., Matthies, H.K., Frey, U., Vorobyev, V.S., and Matthies, H.: Calcium-induced long-term potentiation in the hippocampal slice: Characterization of the time course and conditions. **Brain Res. Bull.**, 1986, 17, 291-296

Pohle, W., Reymann, K.G., Jork, R., and Malisch, R.: The influence of experimental conditions on the morphological preservation of hippocampal slices in vitro. **Biomed. Biochim. Acta**, 1986, 45, 1145-1152

Reymann, K.G., Frey, U., Jork, R., and Matthies, H.: Polymyxin B, an inhibitor of protein kinase C, prevents the maintenance of synaptic long-term potentiation in hippocampal CA1 neurones. **Brain Res.**, 1988, 440, 305-314

Frey, U., Krug, M., Reymann, K.G., and Matthies, H.: Anisomycin, an inhibitor of protein synthesis, blocks late phases of homo- and heterosynaptic long-term potentiation in the hippocampal CA1 region in vitro. **Brain Res.**, 1988, 452, 57-62

Popov, N., Reymann, K.G., Schulzeck, K., Schulzeck, S., and Matthies, H.: Alterations in calmodulin content in fractions of rat hippocampal slices during tetanic and calcium induced long-term potentiation. **Brain Res. Bull.**, 1988, 21, 201-206

Reymann, K.G., Schulzeck, K., Kase, H., and Matthies, H.: Phorbol ester-induced hippocampal long-term potentiation is counteracted by inhibitors of protein kinase C. **Exp. Brain Res.**, 1988, 71, 227-230

Hvalby, O., Reymann, K.G., and Andersen, P.: Intracellular analysis of phorbol ester induced potentiation is counteracted by inhibitors of protein kinase C. **Exp. Brain Res.**, 1988, 71, 588-596

Reymann, K.G., Brödemann, R., Kase, H., and Matthies, H.: Inhibitors of calmodulin and protein kinase C block different phases of hippocampal long-term potentiation. **Brain Res.**, 1988, 461, 388-392

Reymann, K.G., Matthies, H., Schulzeck, K., and Matthies H.: N-methyl-D-aspartate receptor activation is required for the induction of both early and late phases of long-term potentiation in rat hippocampal slices. **Neurosci. Lett.**, 1989, 96, 96-101

Frey, U., Krug, M., Brödemann, R., Reymann, K.G., and Matthies, H.: Long-term potentiation induced in dendrites separated from their CA1 pyramidal somata does not establish a late phase. **Neurosci. Lett.**, 1989, 97, 135-139

Schröder, H., Reymann, K.G., and Matthies, H.: 4beta-phorbol-12, 13-dibutyrate enhances K-stimulated dopamine release from hippocampal slice. **Biomed. Biochem. Acta**, 1989, 48, 425-429

Reymann, K.G., and Matthies, H.: 2-Amino-4-phosphonobutyrate selectively eliminates late phases of long-term potentiation in rat hippocampus. **Neurosci. Lett.**, 1989, 98, 166-171

Schröder, H., and Reymann, K.G.: N-methyl-D-aspartate (NMDA) stimulates the release of dopamine from rat hippocampal slices. **Biomed. Biochim. Acta**, 1990, 41, 281-284

Davies, S.N., Lester, R.A.J., Reymann, K.G., and Collingridge, G.: Temporally distinct pre- and postsynaptic mechanisms maintain long-term potentiation. **Nature**, 1989, 338, 500-503

Collingridge, G.L., Lester, R.A., and Reymann, K.G.: Learning by handshake? **Nature**, 1989, 339, 588

Reymann, K.G., Davies, S.N., Matthies, H., and Collingridge, G.: A K-252b sensitive protein kinase mediates a late postsynaptic phase of long-term potentiation. **Europ. J. Neurosci.**, 1990, 2, 481-486

Krug, M., Jork, R., Reymann, K.G., Wagner, M., and Matthies, H.: The amnesic substance 2-desoxygalactose suppresses the maintenance of hippocampal LTP. **Brain Res.**, 1991, 540, 237-242

Matthies, H., Behnisch, T., Kase, H., Matthies, H.-J., and Reymann, K.G.: Differential effects of protein kinase inhibitors on pre-established long-term potentiation in rat hippocampal neurons in vitro. **Neurosci. Lett.**, 1991, 121, 259-262

Aronica, E., Frey, U., Wagner, M., Schröder, H., Krug, M., Catania, M.V., Nicoletti, F., and Reymann, K.G.: Enhanced sensitivity of metabotropic glutamate receptors after induction of long-term potentiation in rat hippocampus. **J. Neurochem.**, 1991, 57, 376-383

Behnisch, T., Fjodorow, K., and Reymann, K.G.: L-2-amino-3-phosphonopropionate blocks late synaptic long-term potentiation. **NeuroReport**, 1991, 2, 386-388

Frey, U., Matthies, H., Reymann, K.G., and Matthies, H.: The effect of dopaminergic D1 receptor blockade during tetanization on the expression of long-term potentiation in the rat CA1 region in vitro. **Neurosci. Lett.**, 1991, 129, 111-114

Opitz, T., and Reymann, K.G.: Blockade of metabotropic glutamate receptors protects from hypoxic injury in rat CA1 neurons. **NeuroReport**, 1991, 2, 455-457

Angenstein, F., Matthies, H., Staack, S., Reymann, K.G., and Staak, S.: The maintenance of hippocampal long-term potentiation is paralleled by a dopamine-dependent increase in glycoprotein fucosylation. **Neurochem. Int.**, 1992, 21, 403-408

Matthies, H., Brackley, P., Usherwood, P., and Reymann, K.G.: Philanthotoxin-343 blocks LTP but not synaptic transmission in rat hippocampus. **NeuroReport**, 1992, 3, 649-652

Fedorov, N., Screbitsky, V., and Reymann, K.G.: Effects of philanthotoxin-343 on CA1 pyramidal neurons of rat hippocampus in vitro. **Eur. J. Pharmacol.**, 1992, 228, 201-206

Opitz, T., and Reymann, K.G.: (1S, 3R)-ACPD protects synaptic transmission from hypoxia in hippocampal slices. **Neuropharmacol.**, 1993, 32, 103-104

Behnisch, T., and Reymann, K.G.: Co-activation of metabotropic glutamate and N-methyl-D-Aspartate receptors is involved in mechanisms of long-term potentiation maintenance in rat hippocampal CA1 neurons. **Neuroscience**, 1993, 54, 37-47

Matthies, H., and Reymann K.G.: Protein kinase A inhibitors prevent the maintenance of hippocampal long-term potentiation. **NeuroReport**, 1993, 4, 712-714

Fedorov, N.B., and Reymann, K.G.: Simultaneous local pressure microejection of excitatory amino acids and field potential recording with a single micropipette in the hippocampal slice. **J. Neurosci. Meth.**, 1993, 50, 83-90

Riedel, G., and Reymann, K.G.: An antagonist of the metabotropic glutamate receptor prevents LTP in the dentate gyrus of freely moving rats. **Neuropharmacol.**, 1993, 32, 929-931

Sergueeva, O.A., Fedorov, N.B., and Reymann, K.G.: An antagonist of glutamate metabotropic receptors, (RS- α -methyl-4-carboxyphenylglycine, prevents the LTP-related increase in postsynaptic AMPA sensitivity in hippocampal slices. **Neuropharmacol.**, 1993, 32, 933-935

Seidenbecher, T., Balschun, D., Vogel, D., and Reymann, K.G.: Neuronal transmission of hippocampal CA1 neurones is modulated by corticotropin-like intermediate lobe peptide [CLIP; ACTH 18-39]. **Peptides**, 1993, 14, 1221-1224

Behnisch, T., and Reymann, K.G.: 2,3-Diphosphoglyceric acid blocks long-term potentiation of excitatory postsynaptic currents in hippocampal CA1 neurons of the rat. **Neurosci. Lett.**, 1994, 165, 23-26

Riedel, G., Seidenbecher, T., and Reymann, K.G.: LTP in hippocampal CA1 of urethane narcotized rats requires stronger tetanization parameters. **Physiol. Behav.**, **1994**, **55**, 1141-1146

Riedel, G., Wetzel, W., and Reymann, K.G.: (R,S)- α -methyl-4-carboxyphenylglycine (MCPG) blocks spatial learning in rats and LTP in the dentate gyrus in vivo. **Neurosci. Lett.**, **1994**, **167**, 141-144

Angenstein, F., Riedel, G., Reymann, K.G., and Staak, S.: Hippocampal long-term potentiation in vivo induces translocation of protein kinase C γ . **NeuroReport**, **1994**, **5**, 381-384

Opitz T., Richter, P., and Reymann, K.G.: The metabotropic glutamate receptor antagonist (+)- α -methyl-4-carboxyphenylglycine protects hippocampal CA1 neurons of the rat from in vitro hypoxia/hypoglycemia. **Neuropharmacol.**, **1994**, **33**, 715-717

Balschun, D., and Reymann, K.G.: Cholecystinin (CCK-8S) prolongs 'unsaturated' theta-pulse induced potentiation in rat hippocampal CA1 in vitro. **Neuropeptides**, **1994**, **26**, 421-427

Brown, R.E., Rabe, H., and Reymann, K.G.: (RS)- α -methyl-4-carboxyphenylglycine (MCPG) does not block theta burst-induced long-term potentiation in area CA1 of rat hippocampal slices. **Neurosci. Lett.**, **1994**, **170**, 17-21

Wilsch, V.W., Pidoplichko, V.I., Opitz, T., Shinozaki, H., and Reymann, K.G.: DCG IV, a metabotropic glutamate receptor agonist, behaves also as a potent NMDA agonist on acutely isolated rat hippocampal CA1 neurons. **Eur. J. Pharmacol.**, **1994**, **262**, 287-291

Riedel, G., Wetzel, W., and Reymann, K.G.: Computer-assisted shock-reinforced Y-maze training: A method for studying spatial alternation behavior. **NeuroReport**, **1994**, **5**, 2061-2064

Pidoplichko, V.I., and Reymann, K.G.: Abscisic acid potentiates NMDA-gated currents in hippocampal neurons. **NeuroReport**, **1994**, **5**, 2311-2316

Brown, R.E., Fedorov, N.B., Haas, H.L., and Reymann, K.G.: Histaminergic modulation of synaptic plasticity in area CA1 of rat hippocampal slices. **Neuropharmacol.**, **1995**, **34**, 181-190

Riedel, G., Casabona, G., and Reymann, K.G.: Inhibition of long-term potentiation in the dentate gyrus of freely moving rats by the metabotropic glutamate receptor antagonist MCPG. **J. Neurosci.**, **1995**, **15**, 87-98

Seidenbecher, T., Balschun, D., and Reymann, K.G.: Drinking after water deprivation prolongs 'unsaturated' LTP in the dentate gyrus of rats. **Physiol. Behav.**, **1995**, **57**, 1001-1004

Fedorov, N.B., Pasinelli, P., Oestreicher, A.B., DeGraan, P.N.E., and Reymann, K.G.: Antibodies to postsynaptic PKC substrate neurogranin prevent long-term potentiation in hippocampal CA1 neurons. **Eur. J. Neurosci.**, 1995, 7, 819-822

Riedel, G., Wetzel, W. Kozikowski, A.P., and Reymann, K.G.: Block of spatial learning by mGluR agonist tADA in rats. **Neuropharmacol.**, 1995, 34, 559-561

Frey, U., Schollmeier, K., Reymann, K.G., and Seidenbecher, T.: Saturated hippocampal LTP in rats does not preclude additional potentiation at later phases. **Neurosci.**, 1995, 67, 799-807

Opitz, T., Richter, P., Carter, A.J., Kozikowski, A. P., Shinozaki, H., and Reymann, K.G.: Metabotropic glutamate receptor subtypes differentially influence neuronal recovery from in vitro hypoxia/hypoglycemia in rat hippocampal slices. **Neurosci.**, 1995, 68, 989-1001

Manahan-Vaughan, D., and Reymann, K.G.: Regional and developmental profile of modulation of hippocampal synaptic transmission and LTP by AP4-sensitive mGluRs in vivo. **Neuropharmacol.**; 1995, 34, 991-1001

Wilsch, V.W., Pidoplichko, V. I., and Reymann, K.G.: Preferential block of desensitizing AMPA receptor in hippocampal neurons by GAMS. **Europ. J. Neuropharmacol.**, 1995, 284, 141-147

Behnisch, T., and Reymann, K.G.: Thapsigargin blocks long-term potentiation induced by weak, but not strong tetanisation in rat hippocampal CA1 neurons. **Neurosci. Lett.**, 1995, 192, 185-188

Manahan-Vaughan, D., and Reymann, K.G.: 1S,3R-ACPD dose-dependently induces a slow-onset potentiation in the rat hippocampal CA1 region in vivo. **Neuropharmacol.**, 1995, 34, 1103-1105

Riedel, G., Manahan-Vaughan, D., Kozikowski, A.P., and Reymann, K.G.: Metabotropic glutamate receptor agonist trans-Azetidine-2,4-dicarboxylic acid facilitates maintenance of LTP in the dentate gyrus in vivo. **Neuropharmacol.**, 1995, 34, 1107-1109

Platt, B., Carpenter, D.O., Büsselberg, D., Reymann, K.G., and Riedel, G.: Aluminum blocks hippocampal long-term potentiation in rats in vitro and in vivo. **Exp. Neurol.**, 1995, 134, 73-86

Brown, R., and Reymann, K.G.: Effects of glutamate receptor agonists on paired-pulse depression in the dentate gyrus of the rat in vitro. **Neurosci. Lett.**, 1995, 196, 17-20

Wetzel, W., Wagner, T., Reymann, K.G., and Riedel, G.: Open field behavior in rats: role of metabotropic receptors. **NeuroReport**, 1995, 196, 17-20

Manahan-Vaughan, D., and Reymann, K.G.: 1S, 3R-ACPD dose-dependently facilitates NMDA-dependent LTP and induces a slow-onset potentiation in the dentate gyrus in vivo. **Europ. J. Pharmacol.**, 1995, 294, 497-503

Yanovsky, Y., Reymann, K.G., and Haas, H.L.: pH-dependent facilitation of synaptic transmission by histamine in the CA1-region of mouse hippocampus. **Europ. J. Neurosci.**, 1995, 7, 2017-2020

Riedel, G., Wetzel, W., and Reymann, K.G.: Metabotropic glutamate receptors in spatial and nonspatial learning in rats studied by means of agonist and antagonist application. **Learning and Memory**, 1995, 2, 243-265

Platt, B., Behnisch, T., and Reymann, K.G.: Metabotropic glutamate receptors are involved in TEA-induced long-term potentiation in area CA1 of the hippocampus. **Neuropharmacol.**, 1995, 34, 1339-1341

Brown, R.E., and Reymann, K.G.: Class 1 metabotropic glutamate receptor agonists do not facilitate the induction of long-term potentiation in the dentate gyrus of the rat in vitro. **Neurosci. Lett.**, 1995, 202, 73-76

Breakwell, N.A., Behnisch, T., Publicover, S.J., and Reymann, K.G.: Attenuation of high voltage activated Ca²⁺ current run-down in rat hippocampal CA1 pyramidal cells by NaF. **Exp. Brain Res.**, 1995, 106, 505-508

Manahan-Vaughan, D., Reiser, M., Pin, J.-P., Wilsch, V., Kozikowski, A.P., Bockaert, J., Reymann, K.G., and Riedel, G.: Long-term potentiation in rat dentate gyrus in vivo: Different roles for class I metabotropic glutamate receptor subtypes. **Neurosci.**, 1996, 72, 999-1008

Faber-Zuschratter, H., Seidenbecher, T., Reymann, K.G., and Wolf, G.: Ultrastructural distribution of NADPH-diaphorase in the normal hippocampus and after long-term potentiation. **Neurotransmission**, 1996, 103, 807-817

Pidoplichko, V.I., Wilsch, V., and Reymann, K.G.: Fast desensitizing kainate-gated current resolved in whole-cell experiments on isolated hippocampal neurons. **Eur. J. Pharmacol.**, 1996, 303, 145-149

Manahan-Vaughan, D., and Reymann, K.G.: Metabotropic glutamate receptor subtype agonists facilitate long-term potentiation within a distinct time window in the dentate gyrus in vivo. **Neurosci.**, 1996, 74, 723-731

Riedel, G., Opitz, T., and Reymann, K.G.: Blockade of metabotropic glutamate receptors protects hippocampal neurons from hypoxia-induced cell death in rats in vivo. **Progr. Neuro-Psychopharmacol. & Biol. Psychiat.**, 1996, 20, 761-789

Platt, B., and Reymann, K.G.: Inhibition of TEA-induced LTP by Aluminium. **Exp. Neurol.**, 1996, 141, 240-247

Brown, R.E., and Reymann, K.G.: Histamine H3 receptor mediated depression of synaptic transmission in the dentate gyrus of the rat in vitro. **J. Physiol.**, 1996, 496, 175-184

Fedorov, N.B., Brown, R., and Reymann, K.G.: Fast increases of AMPA receptor sensitivity following tetanus-induced potentiation in the CA1 region of the rat hippocampus. **NeuroReport**, 1997, 8, 411-414

Astrelin, A.V., Sokolov, M.V., Behnisch, T., Reymann, K.G., and Voronin, L.L.: Noise deconvolution based on the L1-metric and decomposition of discrete distributions of postsynaptic responses. **J. Neurosci. Meth.**, 1997, 73, 17-27

Seidenbecher, T., Reymann, K.G., and Balschun, D.: A post-tetanic time window for the reinforcement of LTP by appetitive and aversive stimuli. **Proc. Natl. Acad. Sci.**, 1997, 94, 1494-1499

Klein, J., Reymann, K.G., and Riedel, G.: Activation of phospholipases C and D by the novel metabotropic glutamate receptor agonist tADA. **Neuropharmacol.**, 1997, 36, 261-263

Gorkin, A. G., Alexandrov, Y.I., and Reymann, K.G.: Long-term potentiation in cingulate cortex of freely moving rats: duration and mGluR independence. **Neurosc. Res. Commun.**, 1997, 21, 119-124

Manahan-Vaughan, D., and Reymann, K.G.: Group 1 metabotropic glutamate receptors contribute to slow-onset potentiation in the rat CA1 region in vivo. **Neuropharmacol.**, 1998, 36, 1533-1538

Manahan-Vaughan, D., Reymann, K.G., and Brown, R.E.: In vivo electrophysiological investigations into the role of histamine in the dentate gyrus of the rat. **Neurosci.**, 1998, 84, 783-790

Voronin, L.L, Sokolov, M.V., Astrelin, A.V., Behnisch, T., and Reymann, K.G.: Quantal analysis does not confirm increase in sensitivity of postsynaptic receptors during the late phase of hippocampal long-term potentiation. **Dokl. Akad. Nauk**, 1998, 358, 3, 419-421

Sokolov, M.V., Rossokhin, A.V., Behnisch, T., Reymann, K.G., and Voronin, L.L.: Interaction between paired-pulse facilitation and long-term potentiation of minimal excitatory postsynaptic potentials in rat hippocampal slices: a patch clamp study. **Neurosci.**, 1998, 85, 1-13

Astrelin, A.V., Sokolov, M.V., Behnisch, T., Reymann, K.G., and Voronin, L.L.: Principal component analysis of minimal excitatory postsynaptic potentials. **J. Neurosci. Meth.**, 1998, 79, 169-186

Wilsch, V.W., Behnisch, T., Jäger, T., Reymann, K.G., and Balschun, D.: When are class I metabotropic glutamate receptors necessary for long-term potentiation? **J. Neurosci.**, 1998, 18, 6071-6080

Jäger, T., Reymann, K.G., and Behnisch, T.: Analysis of the tetanic and post-tetanic components of intradendritic Ca²⁺ signals in hippocampal CA1 neurons. **Neurosci.**, 1998, 86, 423-429

Behnisch, T., Wilsch, V.W., Balschun, D., and Reymann, K.G.: The role of group II metabotropic glutamate receptors in hippocampal CA1 long-term potentiation in vitro. **Europ. J. Pharmacol.**, 1998, 356, 159-165

Manahan-Vaughan, D., Braunewell K.H., and Reymann, K.G.: Subtype-specific involvement of metabotropic glutamate receptors in two forms of long-term potentiation in the dentate gyrus of freely moving rats. **Neurosci.**, 1998, 86, 709-721

Nakic, M., Manahan-Vaughan, D., Reymann, K.G., and Schachner, M.: Tenascin-C is up-regulated during long-term potentiation in vivo. **J. Neurobiol.**, 1998, 37, 393-404

Bernstein, M., Behnisch, T., Balschun, D., Reymann, K.G., and Reiser, G.: Pharmacological characterisation of metabotropic glutamatergic and purinergic receptors linked to Ca²⁺ signalling in hippocampal astrocytes. **Neuropharmacol.**, 1998, 37, 169-178

Henrich-Noack, P., Hatton, C.D., and Reymann, K.G.: The mGlu receptor ligand (S)-4C3HPG protects neurons after global ischemia in gerbils. **NeuroReport**, 1998, 9, 985-988

Behnisch, T., and Reymann, K.G.: Inhibition of apamin-sensitive calcium dependent potassium channels enhances long-term potentiation in the CA1 region of rat hippocampus in vitro. **Neurosci. Lett.**, 1998, 253, 91-94

Rauca, C., Henrich-Noack, P., Schäfer, K., Höllt, V., and Reymann, K.G.: (S)-4C3HPG reduces infarct size after focal cerebral ischemia. **Neuropharmacol.**, 1998, 37, 1649-1652

Manahan-Vaughan, D., Behnisch, T., Behnisch, G., Vieweg, S., and Reymann, K.G.: Semi-automated analysis of live/ dead neurones in digitised colour images from rat hippocampal thin slices. **J. Neurosci. Meth.**, 1998, 82, 85-95

Schröder, U., Opitz, T., Jäger, T., Sabelhaus, C.F., Breder, J., and Reymann, K.G.: Protective effect of class I metabotropic glutamate receptor activation against hypoxic/hypoglycemic injury in rat hippocampal slices: timing and involvement of PKC. **Neuropharmacol.**, 1999, 38, 209-216

Schröder, U., Opitz, T., Sabelhaus, C.F., Breder, J., and Reymann, K.G.: The novel NA⁺/Ca²⁺ exchange inhibitor KB-R7943 protects CA1 neurons in rat hippocampal slices against hypoxic/hypoglycemic injury. **Neuropharmacol.**, 1999, 38, 319-321

Henrich-Noack, P., and Reymann, K.G.: (1S,3R)-ACPD, a metabotropic glutamate receptor agonist, enhances damage after global ischemia. **Europ. J. Pharmacol.**, 1999, 365, 55-58

Manahan-Vaughan, D., Behnisch, T., and Reymann, K.G.: ACPD-mediated slow-onset potentiation is associated with cell death in the rat CA1 region. **Neuropharmacol.**, 1999, 38, 487-494

Szinyei, C., Behnisch, T., Reiser, G., and Reymann, K.G.: Long-term potentiation in rat CA1 neurons requires post-synaptic calcium induced calcium release. **J. Physiology**, 1999, 516, 855-868

Balschun, D., Manahan-Vaughan, D., Wagner, T., Behnisch, T., Reymann, K.G., and Wetzel, W.: A specific role for mGluRs in hippocampal LTP and hippocampus-dependent spatial learning. **Learning and Memory**, 1999, 6, 138-152

Voronin, L.L., Volgushev, M., Sokolov, M., Kasyanov, A., Chistiakova, M., and Reymann, K.G.: Evidence for an ephaptic feedback in cortical synapses: postsynaptic hyperpolarization alters the number of response failures and quantal content. **Neurosci.**, 1999, 92, 399-405

Angenstein, F., Riedel, G., Reymann, K.G., and Staak, S.: Transient translocation of protein kinase C γ in hippocampal long-term potentiation depends on activation of metabotropic glutamate receptors. **Neurosci.**, 1999, 93, 1289-1295

Manahan-Vaughan, D., Herrero, I., Reymann, K.G., and Sánchez-Prieto, J.: Presynaptic Group 1 metabotropic glutamate receptors may contribute to the expression of long-term potentiation in the hippocampal CA1 region. **Neurosci.**, 1999, 94, 71-82

Kulla, A., Reymann, K.G., and Manahan-Vaughan, D.: Time dependent induction of depotentiation in the dentate gyrus of freely moving rats: involvement of group 2 metabotropic glutamate receptors. **Eur. J. Neurosci.**, 1999, 11, 3864-3872

Henrich-Noack, P., Flor, P.J., Sabelhaus, C., Prass, K., Dirnagl, U., Gasparini, A., Sauter, M., Rudin, M., and Reymann, K.G.: Distinct influence of the group III metabotropic glutamate receptor agonist (R,S)-PPG on different forms of neurodegeneration. **Neuropharmacol.**, 2000, 39, 911-917

Striggow, F., Riek, M., Breder, J., Henrich-Noack, P., Reymann, K.G., and Reiser, G.: The protease thrombin is an endogenous mediator of hippocampal neuroprotection against ischemia at low concentrations but causes degeneration at high concentrations. **Proc. Natl. Acad. Sci.**, 2000, 97, 2264-2269

Breder J., Sabelhaus, C.F., Opitz, T., Reymann, K.G., and Schröder, U.H.: Inhibition of different pathways influencing Na⁺ homeostasis protects organotypic hippocampal slice cultures from hypoxia/hypoglycemia. **Neuropharmacol.**, 2000, 39, 1779-1787

Sabelhaus, C.F., Schröder, U.H., Breder, J., Henrich-Noack, P., and Reymann, K.G.: Neuroprotection against hypoxic/hypoglycaemic injury after the insult by the group III metabotropic glutamate receptor agonist (R,S)-4-phosphonophenylglycine. **Br. J. Pharmacol.**, 2000, 131, 655-658

Kasyanov, A.M., Maximov, V.V., Byzov, A.L., Berretta, N., Sokolov, M.V., Gasparini, S., Cherubini, E., Reymann, K.G., and Voronin, L.L.: Differences in amplitude-voltage relations between minimal and composite mossy fibre responses of rat CA3 hippocampal neurons support the existence of intrasynaptic ephaptic feedback in large synapses. **Neuroscience**, 2000, 101, 323-336

Pak, J.H., Freesia L. Huang, F.L., Li J., Balschun, D., Reymann, K.G., Chiang, C., Westphal, H., and Huang, K.-P.: Neurogranin is involved in the modulation of calcium/calmodulin-dependent protein kinase II, synaptic plasticity, and spatial learning: studies with knockout mice. **Proc. Natl. Acad. Sci.**, 2001, 97, 11232-11237

Striggow, F., Riek-Burchardt, M., Kiesel, A., Schmidt, W., Henrich-Noack, P., Breder, J., Krug, M., Reymann, K.G., and Reiser, G.: Four different types of protease-activated receptors are widely expressed in the brain and are upregulated in hippocampus by severe ischemia. **Eur J. Neurosci.**, 2001, 14, 595-608

Striggow, F., Röhnert, P., and Reymann, K.G.: Telomics: Tissue-based screening for target validation and preclinical drug development. **New Drugs** 1, 2002, 30-36

Wilhelmi, E., Schröder, U., Benabdallah, A., Sieg, F., Breder, J., and Reymann, K.G.: Organotypic brain slice cultures from adult rats: Approaches for a prolonged cultivation time. **ATLA**, 2002, 30, 275-283

Riek-Burchardt, M. Striggow, F., Henrich-Noack, P., Reiser, G., and Reymann, K.G.: Increase of prothrombin-mRNA after global cerebral ischemia in rats, with constant expression of protease nexin-1 and protease-activated receptors. **Neurosci. Lett.**, 2002, 329, 181-184

Schmidt, W., and Reymann, K.G.: Proliferating cells differentiate into neurons in the hippocampal CA1 region of gerbils after global cerebral ischemia. **Neurosci. Lett.**, 2002, 334, 153-156

Jäger, T., Behnisch, T., and Reymann, K.G.: High frequency stimulation-induced dendritic calcium waves in rat hippocampal neurons. **Neurosci. Lett.**, 2002, 335, 103-106

Gorkin, A.G., Reymann, K.G., and Alexandrov, Y.I.: Long-term potentiation and unit evoked responses in the cortex of freely moving rats. **J. Vyss. Nerv. Dejatelnosti**, 2002, 52, 675-685

Manahan-Vaughan, D., Ngomba, R.T., Storto, M., Kulla, A., Catania, M.V., Chiechio, S., Rampello, L., Passarelli, F., Capece, A., Reymann, K.G., and Nicoletti F.: An increased expression of the mGlu5 receptor protein following LTP

induction at the perforant path-dentate gyrus synapse in freely moving rats. **Neuropharmacol.**, 2003, 44, 17-25

Pringle, A.K., Schmidt, W., Deans, J.K. Wulfert, E., Reymann, K.G., and Sundstrom, L.E.: 7-hydroxylated epiandrosterone (7-OH-EPIA) reduces ischaemia-induced neuronal damage both in vivo and in vitro. **Eur. J. Neurosci.**, 2003, 18, 117-124

Rohatgi, T., Henrich-Noack, P., Sedehizade, F., Goertler, M., Wallesch, C.W., Reymann, K.G., and Reiser G.: Transient focal ischemia in rat brain differentially regulates mRNA expression of protease-activated receptors 1 to 4. **Neurosci. Res.**, 2004, 75, 273-279

Riek-Burchardt, M., Henrich-Noack, P., Metz, G.A., and Reymann, K.G.: Detection of chronic sensorimotor impairments in the ladder rung walking task in rats with endothelin-1-induced mild focal ischemia. **J. Neurosci. Meth.**, 2004, 137, 227-233

Nagaraja, R., Schröder, H., Reymann, K.G., and Becker, A.: Group I metabotropic glutamate receptors interfere in different ways with pentylenetetrazole seizures, kindling, and kindling-related learning deficits. **Naunyn-Schmiedeberg's Arch. Pharmacol.**, 2004, 370, 26-34

Martínez-Sánchez, M., Striggow, F., Schröder, U., Kahlert, S., Reymann, K.G., and Reiser, G.: Na⁺ and Ca²⁺ homeostasis pathways, cell death and protection after oxygen-glucose-deprivation in organotypic hippocampal slice cultures. **Neuroscience**, 2004, 128, 729-740

Schröder, U.H., Hock, F.J., Wirth, K., Englert, H.C., and Reymann, K.G.: The ATP-regulated K⁺ -channel inhibitor HMR-1372 affects synaptic plasticity in hippocampal slices. **Eur. J. Pharmacol.**, 2004, 502, 99-104

Huang, K.-P., Huang, F.L., Jäger, T., Li, L., Reymann, K.G., and Balschun, D.: Neurogranin/RC3 enhances long-term potentiation and learning by promoting calcium-mediated signaling. **J. Neurosci.**, 2004, 24, 10660-10669

Laskowski, A., Schmidt, W., Dinkel, K., Martínez-Sánchez, M., and Reymann, K.G.: Growth factors modulate traumatic injury induced neurogenesis in organotypic slice cultures. **Brain Res.**, 2005, 1037, 78-89

Henrich-Noack, P., Gorkin, A.G., Krautwald, K., Pforte, C., Schröder, U.H., and Reymann, K.G.: Tetanus-induced re-activation of evoked spiking in the post-ischemic dentate gyrus. **Neurosci.**, 2005, 133, 571-81

Busse, S., Breder, J., Dinkel, K., Reymann, K.G., and Schröder, U.H.: Inhibitors of cation-chloride-cotransporters affect hypoxic/hypoglycemic injury in hippocampal slices. **Brain Res.**, 2005, 1046, 116-121

Baldauf, K., and Reymann, K.G.: Influence of EGF/bFGF treatment on proliferation, neurogenesis and infarct volume after transient focal ischemia. **Brain Res., 2005, 1056, 158-167**

Chechneva, O., Dinkel, K., Schrader, D., and Reymann, K.G.: Precursor cells from lateral ventricle are present in juvenile organotypic hippocampal slice cultures and display differ from dentate gyrus precursors neurogenic properties. **Neurosci., 2005, 136, 343-355**

Nagaraja, R.Y., Becker, A., Reymann, K.G., and Balschun, D.: Repeated administration of group I mGluR antagonists prevents seizure-induced long-term aberrations in hippocampal synaptic plasticity. **Neuropharmacol., 2005, 49 , 179-187**

Cavaliere, F., Dinkel, K., and Reymann, K.G.: Microglia response and P2 receptor participation in oxygen/glucose deprivation-induced cortical damage. **Neurosci., 2005, 136, 615-623**

Pforte, C., Henrich-Noack, P., Baldauf, K., and Reymann, K.G.: Increase in proliferation and gliogenesis but decrease of early neurogenesis in the rat forebrain shortly after transient global ischemia. **Neurosci., 2005, 136, 1133-1146**

Henrich-Noack, P., Striggow, F., Reiser, G., and Reymann, K.G.: Preconditioning with thrombin can be protective or worsen damage after endothelin-1-induced focal ischemia in rats. **J. Neurosci. Res., 2006, 83, 469-475**

Henrich-Noack, P., Riek-Burchardt, M., Baldauf, K., Reiser, G., and Reymann, K.G.: Focal ischemia induces expression of protease-activated receptor1 (PAR1) and PAR3 on microglia and enhances PAR4 labeling in the penumbra. **Brain Res., 2006, 1070, 232-241**

Neumann, J., Gunzer, M., Gutzeit, H.O., Ullrich, O., Reymann, K.G., and Dinkel, K.: Microglia provide neuroprotection after ischemia. **FASEB J., 2006, 20, 714-716.**

Strokin, M., Chechneva, O., Reymann, K.G., and Reiser, G.: Neuroprotection of rat hippocampal slices exposed to oxygen-glucose deprivation by enrichment with docosahexaenoic acid and by inhibition of hydrolysis of docosahexaenoic acid-containing phospholipids by calcium independent phospholipase A2. **Neurosci., 2006, 140, 547-553**

Chechneva, O., Dinkel, K., Cavaliere, F., Martinez-Sanchez, M., and Reymann, K.G.: Neurogenesis in organotypic hippocampal slice cultures inhibited by neuronal damage and inflammation early after oxygen-glucose deprivation restores at later time point. **Neurobiol. Disease, 2006, 23, 247-259**

Cavaliere, F., Dinkel, K., and Reymann, K.G.: The subventricular zone releases factors which can be protective in oxygen/glucose deprivation-induced cortical damage: an organotypic study. **Exp. Neurol., 2006, 201, 66-74**

Gorba, T., Bradoo, P., Antonic, A., Marvin, K., Xu, D., Lobie, P., Reymann, K.G., Gluckman, P.D., and Sieg, F.: Neural regeneration peptides: A new class of chemoattractive and neuronal survival promoting small, secreted proteins. *Exp. Cell Res.*, 2006, 312, 3060-3074

Braun, H., Bühnemann, C., and Reymann, K.G.: Cultivation of primary cortical neurons from embryonic rats under serum free conditions using Matrigel and StartV-medium. *J. Neurosci. Meth.*, 2006, 157, 32-38

Bühnemann, C., Scholz, A., Bernreuther, C., Malik, C.Y., Braun, H., Schachner, M., Reymann, K.G., (equally contributed with MD), and Dihné, M.: Embryonic stem cell-derived neural precursors survive after transplantation into the ischemic cortex and differentiate into functional neurons. *Brain*, 2006, 129, 3238-48

Reymann, K.G., and Frey, J.U.: The late maintenance of hippocampal LTP: requirements, phases, 'synaptic tagging' 'late associativity' and implications. *Neuropharmacol.*, 2007, 52-24-40

Ziv, Y., Finkelstein, A., Geffen, Y., Kipnis, J., Smirnov, I., Shpilman, S., Vertkin, I., Kimron, M., Lange, A., Hecht, T., Reymann, K.G., Marder, J.B., Schwartz, M., and Yoles, E.: A novel immune-based therapy for stroke induces neuroprotection and supports neurogenesis. *Stroke*, 2007, 38, 774-782

Laskowski, A., Reiser, G., and Reymann, K.G.: Protease-activated receptor-1 induces generation of new microglia in the dentate gyrus of traumatised hippocampal slice cultures. *Neurosci. Lett.*, 2007, 415, 17-21

Riek-Burchardt, M., Henrich-Noack, P., and Reymann, K.G.: No improvement of functional and histological outcome after application of the metabotropic glutamate receptor 5 agonist CHPG in a model of endothelin-1-induced focal ischemia in rats. *Neurosci. Res.*, 2007, 57, 499-503

Baldauf, K., Henrich-Noack, P., and Reymann, K.G.: Detrimental effects of halothane narcosis on damage after Endothelin-1- induced MCAO. *J. Neurosci. Meth.*, 2007, 162, 14-18.

Henrich-Noack, P. Alexandre G. Gorkin, Klaus G. Reymann: Predictive value of changes in electroencephalogram and excitatory postsynaptic field potential for CA1 damage after global ischaemia. *Experimental Brain Research* 2007, 181, 79-86.

Cavaliere, F., Amadio, S., Dinkel, K., Reymann, K., Volonte., C.: P2 receptor antagonist TNP-ATP protects hippocampus from oxygen and glucose deprivation cell death. *J. Pharmacol. Exp. Ther.*, 2007, 323, 70-77.

Henrich-Noack, P., Baldauf, K., Reiser, G., Reymann, K.G.: Pattern of time-dependent reduction of histologically determined infarct volume after focal ischaemia in mice. *Neurosci. Lett.* , 2008, in press